arXiv:0903.1470v1 [math.AT] 9 Mar 2009

THE RATIONAL HOMOTOPY TYPE OF THE SPACE OF SELF-EQUIVALENCES OF A FIBRATION

YVES FÉLIX, GREGORY LUPTON, AND SAMUEL B. SMITH

Abstract. Let $\mathrm{Aut}(p)$ denote the space of all self-fibre-homotopy equivalences of a fibration $p\colon E \to B$. When E and B are simply connected CW complexes with E finite, we identify the rational Samelson Lie algebra of this monoid by means of an isomorphism:

$$\pi_*(\mathrm{Aut}(p)) \otimes \mathbb{Q} \cong H_*(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W)).$$

Here $\wedge V \to \wedge V \otimes \wedge W$ is the Koszul-Sullivan model of the fibration and $\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W)$ is the DG Lie algebra of derivations vanishing on $\wedge V$. We obtain related identifications of the rationalized homotopy groups of fibrewise mapping spaces and of the rationalization of the nilpotent group $\pi_0(\mathrm{Aut}_\sharp(p))$ where $\mathrm{Aut}_\sharp(p)$ is a fibrewise adaptation of the submonoid of maps inducing the identity on homotopy groups.

1. Introduction

Given a fibration $p\colon E \to B$ of connected CW complexes, we study $\mathrm{Aut}(p)$, the space of unpointed homotopy self-equivalences $f\colon E \to E$ satisfying $p \circ f = p$, topologized as a subspace of $\mathrm{Map}(E, E)$. This is a monoid under composition of maps. In general, $\mathrm{Aut}(p)$ is a disconnected space with possibly infinitely many components. The group of components $\pi_0(\mathrm{Aut}(p))$ is the group of fibre-homotopy equivalence classes of self-fibre-homotopy equivalences of p. We denote this group by $\mathcal{E}(p)$. We make a general study of $\mathrm{Aut}(p)$, especially in rational homotopy theory.

The monoid $\mathrm{Aut}(p)$ appears as an object of interest in many different situations. When B is a point, $\mathrm{Aut}(p) \simeq \mathrm{Aut}(E)$ is the monoid of (free) self-homotopy equivalences of the space E, and $\mathcal{E}(p) = \mathcal{E}(E)$, the group of self-equivalences of E. Taking $p\colon PB \to B$ to be the path-space fibration, we have $\mathrm{Aut}(p) \simeq \Omega B$ and $\mathcal{E}(p) = \pi_1(B)$ (see Example 2.4, below). When p is a covering map, $\mathrm{Aut}(p)$ contains the group of deck transformations of the covering.

As regards the homotopy type of $\mathrm{Aut}(p)$, under reasonable hypotheses on p, the monoid $\mathrm{Aut}(p)$ is a grouplike space of CW type (see Proposition 2.2). The path components of a well-pointed grouplike space are all of the same homotopy type. Thus we focus on the path component of the identity map which we denote $\mathrm{Aut}(p)_\circ$. Our first main result gives a complete description of the rational H-homotopy type of this connected grouplike space when E is finite.

Before describing our main results, we review some background and notation. Recall the homotopy groups $\pi_*(G)$ of a connected, grouplike space G admit a natural bilinear pairing called the *Samelson product* of f and g (see [30, Ch.III]). If

---

Date: March 9, 2009.

2000 *Mathematics Subject Classification.* 55P62, 55Q15.

Key words and phrases. fibre-homotopy equivalences, Samelson Lie algebra, function space, Sullivan minimal model, derivation.

G has multiplication μ the pair (G, μ) admits a rationalization as in [17] yielding an H-space $(G_{\mathbb{Q}}, \mu_{\mathbb{Q}})$ which is unique up to H-equivalence. We refer to the H-homotopy type of the pair $(G_{\mathbb{Q}}, \mu_{\mathbb{Q}})$ as the *rational H-type* of (G, μ). If (G, μ) is grouplike then $(G_{\mathbb{Q}}, \mu_{\mathbb{Q}})$ is also. In this case, the rational H-type of G is completely determined by the rational Samelson algebra $(\pi_*(G) \otimes \mathbb{Q}, [\, , \,])$. Specifically, two grouplike spaces are rationally H-equivalent if and only if they have isomorphic Samelson Lie algebras [27, Cor.1]. We identify the rational Samelson Lie algebra of $\mathrm{Aut}(p)_\circ$ in the context of Sullivan's rational homotopy theory. Our reference for rational homotopy theory is [10].

In [29], Sullivan defined a functor $A_{PL}(-)$ from topological spaces to commutative differential graded algebras over $\mathbb{Q}$ (DG algebras for short) that is connected to the cochain algebra functor $C^*(-; \mathbb{Q})$ by a sequence of natural quasi-isomorphisms. Let $\wedge V$ denote the free commutative graded algebra on the graded rational vector space V. A DG algebra (A, d) is a *Sullivan algebra* if $A \cong \wedge V$ and if V admits a basis (v_i) indexed by a well ordered set such that $d(v_i) \in \wedge(v_j, j < i)$. If the differential d has image in the decomposables of $\wedge V$ we say (A, d) is *minimal*. Filtering by product length, a minimal DG algebra is seen to be a Sullivan algebra. A DG algebra (A, d) is a *Sullivan model* for X if (A, d) is a Sullivan algebra and there is a quasi-isomorphism $(A, d) \to A_{PL}(X)$. If (A, d) is minimal it is then the *Sullivan minimal model* of X.

A fibration $p \colon E \to B$ of simply connected CW complexes admits a *relative minimal model* (see [10, Pro.15.6]). This is an injection of DG algebras $I \colon (\wedge V, d) \to (\wedge V \otimes W, D)$ equipped with quasi-isomorphisms η_B and η_E which make the following diagram commutative:

$$\begin{tikzcd}
(\wedge V, d) \arrow[r, "I"] \arrow[d, "\eta_B"', "\simeq"] & (\wedge V \otimes \wedge W, D) \arrow[d, "\eta_E"', "\simeq"] \\
A_{PL}(B) \arrow[r, "A_{PL}(p)"] & A_{PL}(E)
\end{tikzcd} \tag{1}$$

Here $(\wedge V, d)$ is the Sullivan minimal model of B while $(\wedge V \otimes \wedge W, D)$ is a Sullivan (but generally non-minimal) model of E. The differential D satisfies $D(W) \subset (\wedge^+ V \otimes \wedge W) + (\wedge V \otimes \wedge^{\geq 2}(W))$ and further W admits a basis w_i indexed by a well ordered set such that $D(w_i) \in \wedge V \otimes \wedge(w_j, j < i)$.

A derivation θ of degree n of a DG algebra (A, d) will mean a linear map *lowering* degrees by n and satisfying $\theta(ab) = \theta(a)b - (-1)^{n|a|} a\theta(b)$ for $a, b \in A$. We write $\mathrm{Der}^n(A)$ for the vector space of all degree n derivations. The graded vector space $\mathrm{Der}^*(A)$ has the structure of a DG Lie algebra with the commutator bracket $[\theta_1, \theta_2] = \theta_1 \circ \theta_2 - (-1)^{|\theta_1||\theta_2|} \theta_2 \circ \theta_1$ and differential $\mathcal{D}(\theta) = [d, \theta]$. Given a DG subalgebra $B \subseteq A$ we write $\mathrm{Der}^*_B(A)$ for the space of derivations of A that vanish on B. The bracket and differential evidently restrict to give a DG Lie algebra $(\mathrm{Der}^*_B(A), \mathcal{D})$. More generally, given a DG algebra map $\phi \colon A \to A'$ we write $\mathrm{Der}^n(A, A'; \phi)$ for the space of degree n linear maps satisfying $\theta(ab) = \theta(a)\phi(b) - (-1)^{n|a|} \phi(a)\theta(b)$ with differential $\mathcal{D}(\theta) = d_{A'} \circ \theta - (-1)^{|\theta|} \theta \circ d_A$. The pair $(\mathrm{Der}^*(A, A'; \phi), \mathcal{D})$ is then a DG vector space. Given a DG subalgebra $B \subseteq A$ we have the DG vector subspace $(\mathrm{Der}^*_B(A, A'; \phi), \mathcal{D})$ of derivations that vanish on B.

We now describe our main results. Given a fibration $p \colon E \to B$ of simply connected CW complexes, choose and fix a relative minimal model $I \colon (\wedge V, d) \to (\wedge V \otimes \wedge W, D)$ as above. We have:

Theorem 1. *Let $p\colon E \to B$ be a fibration of simply connected CW complexes with E finite. There is an isomorphism of graded Lie algebras*

$$\pi_*(\mathrm{Aut}(p)_\circ) \otimes \mathbb{Q} \;\cong\; H_*(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W)).$$

By the remarks above, Theorem 1 completely determines the rational H-type of $\mathrm{Aut}(p)$. Taking $B = *$, we recover Sullivan's Lie algebra isomorphism

$$\pi_*(\mathrm{Aut}(E)) \otimes \mathbb{Q} \cong H_*(\mathrm{Der}(\wedge W))$$

described in [29, Sec.11]. Here $(\wedge W, d)$ is the minimal model for E.

We obtain Theorem 1 as a sharpened special case of a general calculation of the rational homotopy groups of a fibrewise mapping space. Let $p'\colon E' \to B'$ be a second fibration and $f\colon E' \to E$ be a fibrewise map covering a map $g\colon B' \to B$. Let

$$\mathrm{Map}_g(E', E) = \{h\colon E' \to E \mid p \circ h = g \circ p'\}$$

denote the function space of maps over g topologized as a subspace of $\mathrm{Map}(E', E)$. Write $\mathrm{Map}_g(E', E; f)$ for the path component of f. We compute the rational homotopy groups of this space as an extension of Sullivan's original approach. Fix a relative minimal model $I'\colon (\wedge V', d) \to (\wedge V' \otimes \wedge W', D)$ for $p'\colon E' \to B'$ as above. The map $f\colon E' \to E$ has a model $\mathcal{A}_f\colon \wedge V \otimes \wedge W \to \wedge V' \otimes \wedge W'$. A homotopy class $\alpha\colon S^n \to \mathrm{Map}_g(E', E; f)$ has adjoint $F\colon E' \times S^n \to E$ satisfying $p \circ F(x, s) = g \circ p'(x)$ and $F(x, *) = f(x)$ for all $(x, s) \in E \times S^n$. We show F has a DG model

$$\mathcal{A}_F\colon \wedge V \otimes \wedge W \to \wedge V' \otimes \wedge W' \otimes \left(\wedge(u)/\langle u^2\rangle\right)$$

satisfying $\mathcal{A}_F(\chi) = \mathcal{A}_f(\chi) + u\theta(\chi)$ for $\chi \in \wedge V \otimes \wedge W$. Here $|u| = n$ so that $(\wedge(u)/\langle u^2\rangle, 0)$ is a Sullivan model for S^n. The degree n map θ is then an $\mathcal{A}_f$-derivation as defined above and we show θ vanishes on $\wedge V$, that is, θ is an element of $\mathrm{Der}^n_{\wedge V}(\wedge V \otimes \wedge W, \wedge V' \otimes \wedge W'; \mathcal{A}_f)$. We prove the assignment $\alpha \mapsto \theta$ gives rise to a rational isomorphism:

Theorem 2. *Let $p'\colon E' \to B'$ and $p\colon E \to B$ be fibrations of simply connected CW complexes with E' finite. Let $f\colon E' \to E$ be a fibrewise map with model $\mathcal{A}_f$ as above. There is an isomorphism of vector spaces*

$$\pi_n(\mathrm{Map}_g(E', E; f)) \otimes \mathbb{Q} \;\cong\; H_n(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W, \wedge V' \otimes \wedge W'; \mathcal{A}_f)).$$

for $n \geq 2$. Further, the group $\pi_1(\mathrm{Map}_g(E', E; f))$ is nilpotent and satisfies

$$\mathrm{rank}(\pi_1(\mathrm{Map}_g(E', E; f))) = \dim_{\mathbb{Q}}\left(H_1(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W, \wedge V' \otimes \wedge W'; \mathcal{A}_f))\right).$$

Again, taking $B' = B = *$, we recover known results; in this case the vector space isomorphism

$$\pi_n(\mathrm{Map}(X, Y; f)) \otimes \mathbb{Q} \cong H_n(\mathrm{Der}(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f))$$

for $n \geq 2$ appears in [4, 21, 6] where here $\mathcal{M}_f\colon \mathcal{M}_Y \to \mathcal{M}_X$ is the Sullivan minimal model of $f\colon X \to Y$. The result on $\mathrm{rank}(\pi_1(\mathrm{Map}(X, Y; f)))$ is [22, Th.1]. Our proof of Theorem 2 is an adaptation of the proofs of [21, Th.2.1] and [22, Th.1] with adjustments made for the fibrewise setting.

Finally, as regards the group $\mathcal{E}(p)$ of path components of $\mathrm{Aut}(p)$ we remark that this group is not, generally, a nilpotent group and so not directly amenable to rational homotopy theory. In Section 6, we consider the subgroup $\mathcal{E}_\sharp(p)$ of $\mathcal{E}(p)$ consisting of homotopy equivalence classes of maps $f\colon E \to E$ over B inducing the identity on image and cokernel of the connecting homomorphism in the long exact sequence on homotopy groups of the fibration. The group $\mathcal{E}_\sharp(p)$ is nilpotent and localizes well for E finite by the fibrewise extension of results of Maruyama [23]. Let $D_0 \colon W \to V \oplus W$ be the linear part of D in the Sullivan model for p. Write $W = W_0 \oplus W_1$ where $W_0 = \ker D_0$ and W_1 is a vector space complement. Denote by $\mathrm{Der}^0_\#(\wedge V \otimes \wedge W)$ the vector space of derivations θ of degree 0 of $\wedge V \otimes \wedge W$ that satisfy $\theta(V) = 0$, $\mathcal{D}(\theta) = 0$, $\theta(W_0) \subset V \oplus \wedge^{\geq 2}(V \oplus W)$, and $\theta(W_1) \subset V \oplus W_0 \oplus \wedge^{\geq 2}(V \oplus W)$. Define

$$H_0(\mathrm{Der}_\sharp(\wedge V \otimes \wedge W)) = \mathrm{coker}\,\{\mathcal{D} : \mathrm{Der}^1_{\wedge V}(\wedge V \otimes \wedge W) \to \mathrm{Der}^0_\#(\wedge V \otimes \wedge W)\}.$$

The case $B = *$ in the following result is originally due to Sullivan [29, Sec.11]. See also [26, Th.12].

Theorem 3. *Let $p \colon E \to B$ be a fibration of simply connected CW complexes with E finite. There is a group isomorphism*

$$\mathcal{E}_\#(p)_\mathbb{Q} \cong H_0(\mathrm{Der}_\sharp(\wedge V \otimes \wedge W)).$$

The paper is organized as follows. In Section 2, we prove several general results concerning $\mathrm{Aut}(p)$. We prove Theorem 2 in Section 3 as a consequence of some results in fibrewise DG homotopy theory. We prove Theorem 1 in Section 4 and deduce some consequences in Section 5. Finally, in Section 6 we turn to the group of components $\mathcal{E}(p)$ of $\mathrm{Aut}(p)$ and prove Theorem 3. We conclude with an example to show that $\mathcal{E}(p)$ is not generally a subgroup of $\mathcal{E}(E)$.

2. Basic Homotopy theory of $\mathrm{Aut}(p)$

Function spaces like the monoid $\mathrm{Aut}(p)$ will not generally have the homotopy type of a CW complex, even though the spaces involved in the fibration $p \colon E \to B$ do. When E is finite, however, we can deduce a general result for fibrewise mapping spaces as a consequence of standard results. Suppose given a commutative diagram of connected CW complexes

(2)
$$\begin{tikzcd} E' \arrow[r, "f"] \arrow[d, "p'"'] & E \arrow[d, "p"] \\ B' \arrow[r, "g"] & B \end{tikzcd}$$

with vertical maps fibrations as needed to define the fibrewise mapping space $\mathrm{Map}_g(E', E; f)$ mentioned above. We make use of the fibre sequence

$$\mathrm{Map}_g(E', E; f) \to \mathrm{Map}(E', E; f) \xrightarrow{p_*} \mathrm{Map}(E', B; p \circ f).$$

We say a space X is a *nilpotent* space if X is a connected CW complex with $\pi_1(X)$ nilpotent and acting nilpotently on the higher homotopy groups of X. In this case, X admits a rationalization $\ell_X \colon X \to X_\mathbb{Q}$ [17, Ch.II]. Given a map $f \colon X \to Y$ with Y nilpotent, we write $f_\mathbb{Q} = \ell_Y \circ f \colon X \to Y_\mathbb{Q}$.

Proposition 2.1. *If E' is a finite CW complex and E and B are nilpotent spaces, the components of $\mathrm{Map}_g(E', E)$ are all nilpotent spaces. For each component, composition with $\ell_E \colon E \to E_\mathbb{Q}$ gives a rationalization map:*

$$(\ell_E)_* \colon \mathrm{Map}_g(E', E; f) \to \mathrm{Map}_{g_\mathbb{Q}}(E', E_\mathbb{Q}; f_\mathbb{Q}).$$

Proof. By [25], the function spaces $\mathrm{Map}(E', E; f)$ and $\mathrm{Map}(E', B; p \circ f)$ are of CW type. Thus in the fibre sequence above, the fibre is CW as well by [18, Lem.2.4]. Further, by [17, Cor.II.2.6 and Th.II.3.11], $\mathrm{Map}(E', E; f)$ and $\mathrm{Map}(E', B; p \circ f)$ are nilpotent spaces with rationalizations induced by composition with ℓ_E and ℓ_B. By [17, Th.II.2.2], we obtain $\mathrm{Map}_g(E', E; f)$ is nilpotent and then we see ℓ_E induces a rationalization by the Five Lemma. ☐

As regards the monoid $\mathrm{Aut}(p)_\circ$, we may sharpen the first part of this result:

Proposition 2.2. *Let $p\colon E \to B$ be a fibration of connected CW complexes with connected fibre F. If either E or B is finite then $\mathrm{Aut}(p)$ has the homotopy type of a CW complex and the H-homotopy type of a loop-space.*

Proof. For the CW structure in the case B is finite, we use the identity $\mathrm{Aut}(p) \simeq \Omega\mathrm{Map}(B, BF; h)$ where BF is the base of the universal fibre with fibre $F = p^{-1}(b_0)$ (see [16, Th.1] and [5, Th.3.3]). Since BF is CW in this case, applying [25] again gives the result. In either case, $\mathrm{Aut}(p)$ is a strictly associative CW monoid and so admits a Dold-Lashof classifying space $\mathrm{BAut}(p)$. Thus $\mathrm{Aut}(p) \simeq \Omega\mathrm{BAut}(p)$ [15, Satz.7.3] ☐

By Proposition 2.2, the restriction to E finite is not necessary to rationalize $\mathrm{Aut}(p)_\circ$. However, we will make use of the second statement in Proposition 2.1 in the proof of Theorems 1 and 2.

We next recall an interesting invariant of a connected grouplike space G, the *homotopical nilpotency* of G as studied by Berstein and Ganea [3]. It is defined as follows: Using the homotopy inverse, we have commutator maps $\varphi_n\colon G^n \to G$: Here φ_1 is the identity, $\varphi_2(g,h) = ghg^{-1}h^{-1}$ is the usual commutator, and $\varphi_n = \varphi_2 \circ (\varphi_{n-1} \times \varphi_1)$. The homotopical nilpotency $\mathrm{Hnil}(G)$ is then the least integer n such that φ_{n+1} is null-homotopic. The *rational homotopical nilpotency* $\mathrm{Hnil}_{\mathbb{Q}}(G)$ of $G = (G, \mu)$ is defined to be the homotopical nipotency of $G_{\mathbb{Q}} = (G_{\mathbb{Q}}, \mu_{\mathbb{Q}})$. The inequality $\mathrm{Hnil}_{\mathbb{Q}}(G) \leq \mathrm{Hnil}(G)$ is direct from definitions. When $\mathrm{Hnil}(G) = 1$ (respectively, $\mathrm{Hnil}_{\mathbb{Q}}(G) = 1$) we say G is *homotopy abelian* (respectively, *rational homotopy abelian*).

When $G = \Omega X$ is a loop-space, $\mathrm{Hnil}(G)$ is directly related to the nilpotency $\mathrm{Nil}(\pi_*(G))$ of the Samelson bracket $[\,,\,]$ on $\pi_*(G)$ and to the length $\mathrm{WL}(X)$ of the longest Whitehead bracket in $\pi_*(X)$. If X is a nilpotent space, write $\mathrm{WL}_{\mathbb{Q}}(X)$ for the Whitehead length of the rationalization of $X_{\mathbb{Q}}$ of X.

Proposition 2.3. *Let G be a connected CW loop space, $G \simeq \Omega X$ for some simply connected space X. Then*

$$\mathrm{Hnil}_{\mathbb{Q}}(G) = \mathrm{Nil}(\pi_*(G) \otimes \mathbb{Q}) = \mathrm{WL}_{\mathbb{Q}}(X) \leq \mathrm{WL}(X) = \mathrm{Nil}(\pi_*(G)) \leq \mathrm{Hnil}(G).$$

Proof. The equality $\mathrm{WL}(X) = \mathrm{Nil}(\pi_*(G))$ (and its rationalization) is a consequence of the identification of the Whitehead product with the Samelson product via the isomorphism $\pi_*(\Omega X) \cong \pi_{*+1}(X)$ [30, Th.X.7.10]. The inequality $\mathrm{Nil}(\pi_*(G)) \leq \mathrm{Hnil}(G)$ is [3, Th.4.6]. The equality $\mathrm{Nil}(\pi_*(G) \otimes \mathbb{Q}) = \mathrm{WL}_{\mathbb{Q}}(X)$ is [1, Lem.4.2] (see also [26, Th.3]). Finally, the inequality $\mathrm{WL}(X_{\mathbb{Q}}) \leq \mathrm{WL}(X)$ is immediate from the definition. ☐

We give some examples and direct calculations regarding $\mathrm{Aut}(p)$.

Example 2.4. Let PB be the space of Moore paths at $b_0 \in B$:

$$PB = \{(\omega, r) \,|\, r \geq 0, \omega : (R_{\geq 0}, 0) \to (B, b_0)\,, \text{such that } \omega(s) = \omega(r) \text{ for } r \geq s\,\}\,.$$

Let $p\colon PB \to B$ given by $(\omega, r) = \omega(r)$ be the path-space fibration. Then there is an H-equivalence

$$\mathrm{Aut}(p) \simeq \Omega B$$

where $\Omega B = \{(\sigma, s) \in PB \mid \sigma(s) = b_0\}$ is the space of Moore loops at b_0. To see this, define $\theta\colon \Omega B \to \mathrm{Aut}(p)$ given by $\theta((\sigma, s))(\omega, r) = (\sigma{*}\omega, s{+}r)$ and $\psi\colon \mathrm{Aut}(p) \to \Omega B$. Given $f \in \mathrm{Aut}(p)$, f restricts to an equivalence $f\colon \Omega B \to \Omega B$. Define $\psi(f) = f(c_{b_0}, 0)$ where c_{b_0} is the constant loop. Clearly $\psi \circ \theta = id$. On the other hand, a homotopy $H\colon \mathrm{Aut}(\xi) \times [0, 1] \to \mathrm{Aut}(\xi)$, between $\theta \circ \psi$ and the identity is defined by

$$H(f, t)(\omega, r) = (\omega_{tr}, tr) * f(\omega^b_{tr}, r - tr)\,.$$

Here $\omega^b_s(t) = \omega(t + s)$ and

$$\omega_s(t) = \begin{cases} \omega(t) & t \leq s \\ \omega(s) & t \geq s. \end{cases}$$

Using Proposition 2.3, we conclude

$$\mathrm{Hnil}(\mathrm{Aut}(p)_\circ) \geq \mathrm{WL}(\widetilde{B}) \ \text{ and } \ \mathrm{Hnil}_{\mathbb{Q}}(\mathrm{Aut}(p)_\circ) = \mathrm{WL}_{\mathbb{Q}}(\widetilde{B})$$

where $\widetilde{B}$ denotes the universal cover of B.

Example 2.5. Let $LB = \mathrm{Map}(S^1, B)$ be the free loop space on B. Let $q\colon LB \to B$ the fibration that evaluates a free loop at its base point. By adjointness, we have an H-equivalence

$$\mathrm{Aut}(q) \simeq \Omega\big(\mathrm{Map}(LB, B; q)\big).$$

Now let $p\colon B^{[0,1]} {\to} B$ be evaluation at 1. We then have a map $\theta\colon \Gamma(q; \sigma_0) \to \mathrm{Aut}(p)$ where $\Gamma(q; s_0)$ denotes the space of sections of $q\colon LB \to B$ homotopic to the trivial section $s_0(b) = c_b$. Here we set $\theta(\sigma)(\omega) = \omega * \sigma(\omega(1))$. A proof very similar to that in the preceding example shows that θ is a homotopy equivalence of H-spaces. Now by adjunction $\Gamma(q; \sigma_0)$ is homotopy equivalent to the loop space $\Omega\big(\mathrm{Map}(B, B; 1)\big)$. Therefore we have shown there is a H-equivalence $\mathrm{Aut}(p) \simeq \Omega\big(\mathrm{Map}(B, B; 1)\big)$. Consequently, $\mathrm{Aut}(p)_\circ$ is homotopy abelian since $\mathrm{Map}(B, B; 1)$ is an H-space.

Example 2.6. Observe that when $\pi\colon F \times B \to B$ is the (trivial) product fibration then by adjointness we have

$$\mathrm{Aut}(\pi) \cong \mathrm{Map}(B, \mathrm{Aut}(F))$$

with pointwise multiplication in the latter space. More generally, by the fibre-homotopy invariance of $\mathrm{Aut}(p)$ we have this identification for any fibre-homotopy trivial fibration $p\colon E \to B$ with fibre $F = p^{-1}(b_0)$. By [20, Th.4.10], we have

$$\mathrm{Hnil}(\mathrm{Aut}(p)_\circ) = \mathrm{Hnil}(\mathrm{Aut}(F)_\circ)$$

in this case.

Example 2.7. We next consider a class of fibrations of particular interest in rational homotopy theory. An *F_0-space* F is a simply connected complex with $H^*(F; \mathbb{Q})$ and $\pi_*(F) \otimes \mathbb{Q}$ both finite-dimensional and $H^{odd}(F; \mathbb{Q}) = 0$. Regarding these spaces, we have the following famous conjecture:

Conjecture 2.8. (Halperin [14]) *The rational Serre spectral sequence collapses at the* E_2*-term for every fibration* $p\colon E \to B$ *of simply connected CW complexes with fibre* F *an* F_0*-space.*

A connection to the homotopical nilpotency of self-equivalences is provided by a theorem due to Thomas and Meier: An F_0-space F satisfies the Conjecture 2.8 if and only if $\mathrm{Aut}(F)_\circ$ has vanishing even degree rational homotopy groups (see [24, Th.A]). Thus $\mathrm{Nil}(\pi_*(\mathrm{Aut}(F)_\circ) \otimes \mathbb{Q}) = 1$ in this case for degree reasons and so $\mathrm{Hnil}_{\mathbb{Q}}(\mathrm{Aut}(F)_\circ) = 1$. By Example 2.6, we conclude that $\mathrm{Aut}(p)_\circ$ is rationally homotopy abelian for any trivial fibration $p\colon E \to B$ of simply connected spaces with fibre F an F_0-space satisfying Conjecture 2.8.

We extend this calculation from the trivial fibration to any fibration using the argument given in [12, Th.2.3] and [19, Th.5.6] in the gauge group setting. Let F be an F_0-space satisfying Conjecture 2.8. Let $p\colon E \to B$ be any fibration of simply connected CW complexes with B finite. By Gottlieb's identity mentioned above, we have $\mathrm{Aut}(p)_\circ \simeq \Omega_\circ \mathrm{Map}(B, \mathrm{BAut}_1(F); h)$. Here we write $\mathrm{BAut}_1(F) = \mathrm{BAut}(F)_\circ$ for the classifying space of the connected monoid $\mathrm{Aut}(F)_\circ$. By Meier's result, $\mathrm{Aut}(F)_\circ$ has oddly graded rational homotopy groups. Thus $\mathrm{BAut}_1(F)$ has evenly graded rational homotopy groups and so $\mathrm{BAut}_1(F)$ is a rational grouplike space. We thus have an equivalence $\mathrm{Map}(B, \mathrm{BAut}_1(F)_{\mathbb{Q}}; h_{\mathbb{Q}}) \simeq \mathrm{Map}(B, \mathrm{BAut}_1(F)_{\mathbb{Q}}; 0)$ which loops to an H-equivalence $\Omega_\circ \mathrm{Map}(B, \mathrm{BAut}_1(F)_{\mathbb{Q}}; h_{\mathbb{Q}}) \simeq \Omega_\circ \mathrm{Map}(B, \mathrm{BAut}_1(F)_{\mathbb{Q}}; 0)$. Combining, we see

$$\begin{aligned} (\mathrm{Aut}(p)_\circ)_{\mathbb{Q}} &\simeq \Omega_\circ \mathrm{Map}(B, \mathrm{BAut}_1(F)_{\mathbb{Q}}; 0) \quad \simeq \mathrm{Map}(B, \Omega_\circ \mathrm{BAut}_1(F)_{\mathbb{Q}}; 0) \\ &\simeq \mathrm{Map}(B, (\mathrm{Aut}(F)_\circ)_{\mathbb{Q}}; 0). \end{aligned}$$

Thus $\mathrm{Aut}(p)_\circ$ is rationally homotopically abelian since $\mathrm{Aut}(F)_\circ$ as in Example 2.6. We remark that Halperin's Conjecture has been confirmed in many special cases including the case $F = G/H$ where G is any compact Lie group and H is a closed subgroup of maximal rank [28].

Example 2.9. Let $p\colon E \to B$ be a principal G-bundle. Multiplication by an element of G induces a morphism of H-spaces $G \to \mathrm{Aut}(p)$ while evaluation at the identity gives a left inverse $\mathrm{Aut}(p) \to G$. Thus G is a retract of $\mathrm{Aut}(p)$. It follows easily that

$$\mathrm{Hnil}(\mathrm{Aut}(p)_\circ) \geq \mathrm{Hnil}(G_\circ).$$

3. Rational homotopy groups of fibrewise mapping spaces

We now consider the diagram of fibrations (2) above defining the fibrewise mapping space $\mathrm{Map}_g(E, E'; f)$. We assume all the spaces E', E, B', B are simply connected CW complexes. Our calculation of $\pi_n(\mathrm{Map}_g(E', E; f)) \otimes \mathbb{Q}$ will follow the line of proof of [21, Th.2.1]. Namely, we will define a homomorphism Φ' from the ordinary homotopy group of the function space to the DG vector space of derivations and prove Φ' induces an isomorphism Φ after rationalization. As in [21], the construction depends on some DG algebra homotopy theory

We first recall the construction of the relative minimal model of the fibration $p\colon E \to B$ as in the diagram (1) above. Writing $A_{PL}(X) = (A(X), \delta_X)$ for the DG algebra of Sullivan polynomial forms, the relative model of the map $A(p)\colon (A(B), \delta_B) \to (A(E), \delta_E)$ is of the form $(A(B) \otimes \wedge W, D)$ [10, Sec.14]. Choose a minimal model $(\wedge V, d) \to (A(B), \delta_B)$ (the Sullivan minimal model of B) and obtain a quasi-isomorphism

$$\eta_E \colon (\wedge V \otimes \wedge W, D) \to (A(B) \otimes \wedge W), \delta).$$

We take $(A(E), \delta_E) = (A(B) \otimes \wedge W, \delta)$.

Next consider the fibrewise map $f \colon E' \to E$ over $g \colon B' \to B$. We claim a model

$$\mathcal{A}_f \colon (\wedge V \otimes \wedge W, D) \to (\wedge V' \otimes \wedge W', D')$$

for $f \colon E' \to E$ may be constructed as a map "under" $\mathcal{M}_g \colon (\wedge V, d) \to (\wedge V', d')$, the minimal model of g. That is, $\mathcal{A}_f(\chi) = I' \circ \mathcal{M}_g(\chi)$ for $\chi \in \wedge V$ where $I' \colon \wedge V' \to \wedge V' \otimes \wedge W'$ is the inclusion. The proof is direct: we simply invoke the relative lifting lemma [10, Lem.14.4] when lifting in the standard construction of a Sullivan model of a map [10, Sec.12(c)]. Since we will need the notation, we give the details.

Recall the standard construction depends on the *surjective trick*: Given a graded algebra U we define $(S(U), \delta)$ to be the contractible DG Lie algebra on a basis $U \oplus \delta(U)$. Given a DG algebra map $\eta \colon (B, d) \to (A, d)$, this manoeuvre results in a diagram

$$\begin{tikzcd}
B \arrow[r, shift left, "\alpha"] \arrow[d, "\eta"'] & B \otimes S(A) \arrow[l, shift left, "\beta"] \arrow[dl, two heads, "\gamma"] \\
A &
\end{tikzcd}$$

in which γ is a surjection, and both α and β are quasi-isomorphisms. Write $\eta_B \colon (\wedge V, d) \to (A(B), \delta_B)$ and $\eta_{B'} \colon (\wedge V', d') \to (A(B'), \delta_{B'})$ for Sullivan mininal models for B and B'. Convert $\eta_{B'}$ into a surjection $\gamma_{B'} \colon \wedge V' \otimes S(A(B')) \to A(B')$ as above. We can then lift the composite $A(g) \circ \eta_B$ through the surjective quasi-isomorphism $\gamma_{B'}$, using the standard lifting lemma [10, Lem.12.4]. We thus obtain $\phi_g \colon \wedge V \to \wedge V' \otimes S(A(B'))$. Now set $\mathcal{M}_g = \beta_{B'} \circ \phi_g$. All this is summarized in the following diagram.

$$\begin{tikzcd}
 & & \wedge V' \otimes S(A(B')) \arrow[dl, shift left, "\beta_{B'}", "\simeq"'] \arrow[ddl, bend left, two heads, "\gamma_{B'}", "\simeq"'] \\
\wedge V \arrow[r, dotted, "\mathcal{M}_g"'] \arrow[d, "\eta_B"', "\simeq"] \arrow[urr, bend left, "\phi_g"] & \wedge V' \arrow[ur, shift left, "\alpha_{B'}", "\simeq"'] \arrow[d, "\eta_{B'}"', "\simeq"] & \\
A(B) \arrow[r, "A(g)"'] & A(B') &
\end{tikzcd}$$

Here the symbol $\simeq$ indicates a map is a quasi-isomorphism. By construction, we have $\gamma_{B'} \circ \phi_g = A(g) \circ \eta_B$.

We now apply the same construction to obtain a model $\mathcal{A}_f$ as above. That is, we obtain a lifting ϕ_f in the diagram

$$\begin{tikzcd}
\wedge V \otimes \wedge W \arrow[r, dotted, "\phi_f"] \arrow[dr, "A(f) \circ \eta_E"'] & \wedge V' \otimes \wedge W' \otimes S(A(E')) \arrow[d, two heads, "\gamma_{E'}", "\simeq"'] \\
 & A(E')
\end{tikzcd}$$

and set $\mathcal{A}_f = \beta_{E'} \circ \phi_f$. However, in this case we invoke [10, Lem.14.4] and choose the lifting ϕ_f relative to the partial lifting $\phi'_g\colon \wedge V \to \wedge V' \otimes \wedge W' \otimes S(A(E'))$, where $\phi'_g = S(A(p')) \circ \phi_g$ and $S(A(p'))\colon S(A(B')) \to S(A(B))$ is the map induced by $A(p')\colon A(B) \to A(B')$. The result is the DG model $\mathcal{A}_f$ under $\mathcal{M}_g$.

Now let $\alpha \in \pi_n(\mathrm{Map}_g(E, E'; f))$ be a homotopy element with adjoint $F\colon E' \times S^n \to E$. Write $i\colon E' \to E' \times S^n$ for the based inclusion and $q\colon E' \times S^n \to B'$ for p' composed with the projection. The map F is then a map "under" f and "over" g. That is, $F \circ i = f$ and $p \circ F = g \circ q$. The definition of $\Psi'(\alpha)$ requires a particular form for the Sullivan model of F: We need that F admits a Sullivan model which is "over" $\mathcal{A}_f$ and "under" $\mathcal{M}_g$ in the appropriate sense. Further, we will need the corresponding result for homotopies of F and for the adjoint of a sum of two homotopy classes. We also need a converse result giving spatial realization. Our needs are covered by the following two results.

Let Z be a nilpotent space. The diagram (2) gives rise to the diagram:

(3)
$$\begin{tikzcd}
E' \arrow[d, "i"'] \arrow[dr, "f"] & \\
E' \times Z \arrow[r, dotted, "F"] \arrow[d, "q"'] & E \arrow[d, "p"] \\
B' \arrow[r, "g"] & B
\end{tikzcd}$$

We are interested in maps F making the diagram strictly commute.

Let $\eta_Z\colon (C, d) \to (A(Z), \delta_Z)$ be a Sullivan model for Z. On the DG algebra side, we have a corresponding diagram:

(4)
$$\begin{tikzcd}
\wedge V \arrow[r, "\mathcal{M}_g"] \arrow[d, "I"'] & \wedge V' \arrow[d, "I''"] \\
\wedge V \otimes \wedge W \arrow[r, dotted, "\Gamma"] \arrow[dr, "\mathcal{A}_f"'] & \wedge V' \otimes \wedge W' \otimes C \arrow[d, "P"] \\
& \wedge V' \otimes \wedge W'.
\end{tikzcd}$$

Here I'' is the inclusion induced by I' and P is the obvious projection. Again we consider Γ making the diagram strictly commute. Recall we are assuming the spaces E, E', B, B' are all simply connected CW complexes.

Proposition 3.1. *A Sullivan model $\mathcal{A}_F$ for a map F making (3) commute may be chosen so that $\Gamma = \mathcal{A}_F$ makes (4) commute.*

Proof. The existence of a model $\mathcal{A}_F$ satisfying $P \circ \mathcal{A}_F = \mathcal{A}_f$ is given by [21, Pro.A.2]. Precisely, there is a lifting ϕ_F of $\eta_E \circ A(F)$ as shown:

$$\begin{tikzcd}
\wedge V \otimes \wedge W \arrow[r, dotted, "\phi_F"] \arrow[dr, "\eta_E \circ A(F)"'] & \wedge V' \otimes \wedge W' \otimes C \otimes S(A(E' \times Z)) \arrow[d, "\lambda_{E' \times Z}"] \\
& A(E' \times Z)
\end{tikzcd}$$

which is compatible with the given lift ϕ_f for $\eta_E \circ A(f)$ constructed above. As above, we may obtain the lift ϕ_F as an extension of the partial lift

$$\phi'_g\colon \wedge V \to \wedge V' \otimes \wedge W' \otimes C \otimes S(A(E' \times Z))$$

induced by the map $\phi_g\colon \wedge V \to \wedge V' \otimes \wedge W' \otimes S(A(B'))$ constructed above.

We note that [21, Pro.A.2] is stated in terms of Sullivan minimal models. However, Sullivan models are sufficient: results used there (e.g., [21, Pro.A.4]) require only that the domains of lifts be Sullivan algebras. $\square$

Proposition 3.2. *Suppose E' and Z are finite CW complexes. Let Γ be given making (4) commute. Then Γ is the Sullivan model for a map $F_0\colon E' \times Z \to E_{\mathbb{Q}}$ making the diagram*

$$\begin{tikzcd}
E' \arrow[d, "i"'] \arrow[dr, "f_{\mathbb{Q}}"] & \\
E' \times Z \arrow[r, "F_0"] \arrow[d, "q"'] & E_{\mathbb{Q}} \arrow[d, "p_{\mathbb{Q}}"] \\
B' \arrow[r, "g_{\mathbb{Q}}"] & B_{\mathbb{Q}}
\end{tikzcd}$$

commute.

Proof. First assume $p_{\mathbb{Q}}\colon E_{\mathbb{Q}} \to B_{\mathbb{Q}}$ is a principal $K(W_n, n)$-fibration for W_n a rational space concentrated in degree n. We then have a map $P\colon E_{\mathbb{Q}} \times K(W_n, n) \to E_{\mathbb{Q}}$, the fibrewise action, which is a map over 1_B. By hypothesis, Γ is of the form $\Gamma(\chi) = \mathcal{A}_f(\chi) + \eta(\chi)$ where $\eta(\chi) = 0$ for $\chi \in \wedge V$ and $\eta(w)$ is in the ideal generated by C^+ in $\wedge V' \otimes \wedge W' \otimes C$ for $w \in W_n$. It follows that each $\eta(w)$ is a cycle. By the standard correspondence between cohomology classes and maps into $K(W_n, n)$, we obtain a map $\gamma\colon E' \times Z \to K(W_n, n)$ induced by η. Let $\widetilde{f_{\mathbb{Q}}} = f_{\mathbb{Q}} \circ \pi\colon E' \times Z \to E$ where $\pi\colon E' \times Z \to E'$ is the projection. Define F_0 to be the composite

$$E' \times Z \xrightarrow{\Delta} E' \times Z \times E' \times Z \xrightarrow{\widetilde{f_{\mathbb{Q}}} \times \gamma} E_{\mathbb{Q}} \times K(W_n, n) \xrightarrow{P} E_{\mathbb{Q}}$$

where Δ is the diagonal. It is direct to check F_0 has Sullivan model Γ.

Now proceed by induction over a Moore-Postnikov factorization of the fibration $p_{\mathbb{Q}}\colon E_{\mathbb{Q}} \to B_{\mathbb{Q}}$. Let $p_n\colon (E_{\mathbb{Q}})_n \to (E_{\mathbb{Q}})_{n-1}$ be the nth fibration, a principal fibration with fibre $K(W_n, n)$. Let $\Gamma_n\colon \wedge V \otimes \wedge W_{(n)} \to \wedge V' \otimes \wedge W' \otimes C$ be the restriction of Γ where $W_{(n)} = \bigoplus_{k \leq n} W_k$. By induction, we obtain a map $F_n\colon E' \times Z \to (E_n)_{\mathbb{Q}}$ making the corresponding diagram (3) commute. Since $E' \times Z$ is finite, for n large, composition with the canonical map $h_n\colon E_{\mathbb{Q}} \to (E_{\mathbb{Q}})_n$ yields a bijection $[E' \times Z, E_{\mathbb{Q}}] \cong [E' \times Z, (E_{\mathbb{Q}})_n]$. We choose $F_0\colon E' \times Z \to E_{\mathbb{Q}}$ with $F_0 = F_n \circ h_n$ for such an n. $\square$

Again let $\alpha \in \pi_n(\mathrm{Map}_g(E', E; f))$ with adjoint $F\colon E' \times S^n \to E$ be given. By Proposition 3.1, F has a Sullivan model

$$\mathcal{A}_F\colon \wedge V \otimes \wedge W \to \wedge V' \otimes \wedge W' \otimes (\wedge(u)/\langle u^2 \rangle) \quad \text{with} \quad \mathcal{A}_F(\chi) = \mathcal{A}_f(\chi) + \theta(\chi)u$$

for $\chi \in \wedge V \otimes \wedge W$ where $\mathcal{A}_f(\chi) = \mathcal{M}_g(\chi)$ and $\theta(\chi) = 0$ for $\chi \in \wedge V$. Here $(\wedge(u)/\langle u^2 \rangle, 0)$ is a Sullivan model for S^n with $|u| = n$. The map θ is thus linear of degree n vanishing on $\wedge V$. The following facts are standard and direct to check:

(1) $\mathcal{A}_F(\chi_1\chi_2) = \mathcal{A}_F(\chi_1)\mathcal{A}_F(\chi_2) \Longrightarrow \quad \theta$ is an $\mathcal{A}_f$-derivation.

(2) $\mathcal{A}_F \circ D = D \circ \mathcal{A}_F \Longrightarrow \quad \theta$ is an $\mathcal{A}_f$-derivation cycle.

Define

$$\Phi'\colon \pi_n(\mathrm{Map}_g(E', E; f)) \to H_n(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W, \wedge V' \otimes \wedge W'; \mathcal{A}_f))$$

by $\Phi'(\alpha) = \langle\theta\rangle$. The following result contains the first assertion of Theorem 2.

Theorem 3.3. *Suppose given a commutative diagram*

$$\begin{tikzcd} E' \arrow[r, "f"] \arrow[d, "p'"'] & E \arrow[d, "p"] \\ B' \arrow[r, "g"] & B \end{tikzcd}$$

with vertical maps fibrations and all spaces simply connected CW complexes. The map

$$\Phi'\colon \pi_n(\mathrm{Map}_g(E', E; f)) \to H_n(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W, \wedge V' \otimes \wedge W'; \mathcal{A}_f))$$

defined above is a well-defined map of sets for $n \geq 1$ and a homomorphism for $n \geq 2$. If E' is finite, the rationalization

$$\Phi\colon \pi_n(\mathrm{Map}_g(E', E; f)) \otimes \mathbb{Q} \longrightarrow H_n(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W, \wedge V' \otimes \wedge W'; \mathcal{A}_f))$$

of Φ' is an isomorphism for $n \geq 2$ and a surjection of sets for $n = 1$.

Proof. We prove the first two assertions by adapting the corresponding proofs in [21, Th.2.1] except here we use Proposition 3.1 to model adjoint maps and homotopies. We first show Φ is well-defined for $n \geq 1$. Observe a homotopy $\alpha \sim \beta$ between classes in $\pi_n(\mathrm{Map}_g(E', E; f))$ has adjoint $H\colon E' \times S^n \times I \to E$ giving rise to a commutative diagram (3) with $Z = S^n \times I$. Writing $\wedge(t, dt)$ for the standard model for $I = [0, 1]$ with $|t| = 0$, by Proposition 3.1, H has a Sullivan model

$$\mathcal{H}\colon\ \wedge V \otimes \wedge W \to \wedge V' \otimes \wedge W' \otimes (\wedge(u)/\langle u^2\rangle) \otimes \wedge(t, dt)$$

taking the form

$$\mathcal{H}(\chi) = \mathcal{A}_f(\chi) + \theta_a(\chi)u + (\theta_b(\chi) - \theta_a(\chi))tu + \overline{\theta}(\chi)dt + \text{ terms involving } t^2 \text{ and } tdt.$$

Here $\theta_a, \theta_b \in \mathrm{Der}^n_{\wedge V}(\wedge V \otimes \wedge W, \wedge V' \otimes \wedge W')$ are cycles representing $\Phi'(\alpha)$ and $\Phi'(\beta)$. The map $\overline{\theta}$ is directly checked to be an $\mathcal{A}_f$-derivation of degree $n+1$ vanishing on $\wedge V$ and satisfying $\mathcal{D}(\overline{\theta}) = \theta_b - \theta_a$.

To show that Φ' is a homomorphism for $n \geq 2$, let $\alpha, \beta \in \pi_n(\mathrm{Map}_g(E', E; f))$ with adjoints $F, G\colon E' \times S^n \to E$. Let $(F \mid G)\colon E' \times (S^n \vee S^n) \to E$ be the map induced by F and G. We then have a commutative diagram (3) with $Z = S^n \vee S^n$. A Sullivan model for $S^n \vee S^n$ is $(\wedge(u, v)/\langle u^2, uv, v^2\rangle, 0)$ with $|u| = |v| = n$. Applying Proposition 3.1, we see $(F \mid G)$ has Sullivan model

$$\chi \mapsto \chi + \theta_a(\chi)u + \theta_b(\chi)v\colon\ \wedge V \otimes \wedge W \to \wedge V' \otimes \wedge W' \otimes (\wedge(u, v)/\langle u^2, uv, v^2\rangle)$$

for $\chi \in \wedge V \otimes \wedge W$. Using that $(F \mid G) \circ i_1 = F$ and $(F \mid G) \circ i_2 = G$, where $i_j\colon S^n \to S^n \vee S^n$ are the inclusions, gives θ_a and θ_b are cycle representatives for $\Phi'(\alpha)$ and $\Phi'(\beta)$, respectively. The map $(1_{E'} \times \sigma) \circ (F \mid G)$ is adjoint to the sum

$\alpha+\beta$ where $\sigma\colon S^n \to S^n \vee S^n$ is the pinch map. The result now follows from the fact that a Sullivan model for σ is given by

$$u, v \mapsto w\colon \wedge(u,v)/\langle u^2, uv, v^2\rangle \to \wedge(w)/\langle w^2\rangle.$$

Now assume E' is finite. By Propositon 2.1, composition with a rationalization $\ell_E\colon E \to E_{\mathbb{Q}}$ gives a rationalization $\ell_E\colon \mathrm{Map}_g(E', E; f) \to \mathrm{Map}_{g_{\mathbb{Q}}}(E', E_{\mathbb{Q}}; f_{\mathbb{Q}})$. We take

$$\Phi\colon \pi_n(\mathrm{Map}_{g_{\mathbb{Q}}}(E', E_{\mathbb{Q}}; f_{\mathbb{Q}})) \to H_n(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W, \wedge V' \otimes \wedge W'; \mathcal{A}_f))$$

to be the map Φ' corresponding to the diagram

$$\begin{tikzcd} E' \arrow[r, "f_{\mathbb{Q}}"] \arrow[d, "p'"'] & E_{\mathbb{Q}} \arrow[d, "p_{\mathbb{Q}}"] \\ B' \arrow[r, "g_{\mathbb{Q}}"] & B_{\mathbb{Q}} \end{tikzcd}.$$

Suppose $\Phi(\alpha) = 0$ for $\alpha \in \pi_n(\mathrm{Map}_{g_{\mathbb{Q}}}(E', E_{\mathbb{Q}}; f_{\mathbb{Q}}))$. Then by Proposition 3.1, a Sullivan model $\mathcal{A}_F$ for the adjoint $F\colon E' \times S^n \to E_{\mathbb{Q}}$ is given by:

$$\chi \mapsto \mathcal{A}_f(\chi) + \theta(\chi)u\colon \wedge V \otimes \wedge W \to \wedge V' \otimes \wedge W' \otimes (\wedge(u)/\langle u^2\rangle)$$

and $\theta = \mathcal{D}(\overline{\theta})$ for some $\overline{\theta} \in \mathrm{Der}^{n+1}_{\wedge V}(\wedge V \otimes \wedge W, \wedge V' \otimes \wedge W'; \mathcal{A}_f)$. Define a DG algebra homotopy $\mathcal{H}$ from the map

$$\chi \mapsto \mathcal{A}_f(\chi)\colon \wedge V \otimes \wedge W \to \wedge V' \otimes \wedge W' \otimes (\wedge(u)/\langle u^2\rangle)$$

to the map $\mathcal{A}_F$ by the rule

$$\chi \mapsto \mathcal{A}_f(\chi) + \theta(\chi)ut + \overline{\theta}(\chi)dt\colon \wedge V \otimes \wedge W \to \wedge V' \otimes \wedge W' \otimes (\wedge(u)/\langle u^2\rangle) \otimes \wedge(t, dt).$$

Using Proposition 3.2 with $Z = S^n \times I$, we obtain a homotopy $H\colon E' \times S^n \times I \to E_{\mathbb{Q}}$ between the adjoint of the trivial class in $\pi_n(\mathrm{Map}_{g_{\mathbb{Q}}}(E', E_{\mathbb{Q}}; f_{\mathbb{Q}}))$ and F.

Finally, to prove Φ is onto for $n \geq 1$, let $\theta \in \mathrm{Der}^n_{\wedge V}(\wedge V \otimes \wedge W, \wedge V' \otimes \wedge W'; \mathcal{A}_f)$ be a cycle. Define a DG algebra map Γ by

$$\chi \mapsto \mathcal{A}_f(\chi) + u\theta(\chi)\colon \wedge V \otimes \wedge W \to \wedge V' \otimes \wedge W' \otimes C.$$

Using Proposition 3.2 with $Z = S^n$, we obtain the adjoint $F\colon E' \times S^n \to E_{\mathbb{Q}}$ to a class $\alpha \in \pi_n(\mathrm{Map}_{g_{\mathbb{Q}}}(E', E_{\mathbb{Q}}; f_{\mathbb{Q}}))$ with $\Phi(\alpha) = \langle\theta\rangle$. □

We note that $\pi_1(\mathrm{Map}_g(E', E; f))$ is not, in general, abelian and so Φ cannot, in general, be an isomorphism (c.f. [22, Ex.1.1]). The proof that Φ' is a homomorphism breaks down because $Z = S^1 \vee S^1$ is a non-nilpotent space. In fact, Φ is generally not a homomorphism in degree 1.

By Proposition 2.1, $\pi_1(\mathrm{Map}_g(E', E; f)$ is a nilpotent group though and thus has a well-defined rank. We have:

Theorem 3.4. *With hypotheses as in Theorem 3.3, if E' is finite then*

$$\mathrm{rank}(\pi_1(\mathrm{Map}_g(E', E; f))) = \dim(H_1(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W, \wedge V' \otimes \wedge W'; \mathcal{A}_f)).$$

Proof. The proof is an adaptation of [22, Th.1] similar to the preceding result. Here we use a Moore-Postnikov factorization of $p_{\mathbb{Q}}\colon E_{\mathbb{Q}} \to B_{\mathbb{Q}}$ in place of the absolute Postnikov tower of Y for $\mathrm{Map}(X, Y; f)$ used there. Also we use Theorem 3.3 in place of [21, Th.2.1]. We omit the details. □

4. The rational Samelson Lie Algebra of $\mathrm{Aut}(p)_\circ$

In this section, we sharpen Theorem 3.3 in the case f and g are the respective identity maps to prove Theorem 1. Fix a fibration $p\colon E \to B$ of simply connected CW complexes with E finite. Observe $\mathrm{Aut}(p)_\circ = \mathrm{Map}_{1_B}(E, E; 1_E)$. We prove the map

$$\Phi'\colon \pi_*(\mathrm{Aut}(p)_\circ) \to H_*(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W))$$

constructed above is an isomorphism in degree $n = 1$ and induces an isomorphism Φ of graded Lie algebras after rationalization of the domain.

Let $\alpha \in \pi_p(\mathrm{Aut}(p)_\circ)$ and $\beta \in \pi_q(\mathrm{Aut}(p)_\circ)$ be homotopy classes with adjoints $F\colon E \times S^p \to E$ and $G\colon E \times S^q \to E$. Let $\theta_a \in \mathrm{Der}^p_{\wedge V}(\wedge V \otimes \wedge W)$ and $\theta_b \in \mathrm{Der}^q_{\wedge V}(\wedge V \otimes \wedge W)$ be cycle representatives for $\Phi'(\alpha)$ and $\Phi'(\beta)$, respectively. Define a homotopy class $\alpha * \beta \in [S^p \times S^q, \mathrm{Aut}(p)_\circ]$ to be the composite

$$S^p \times S^q \xrightarrow{\alpha \times \beta} \mathrm{Aut}(p)_\circ \times \mathrm{Aut}(p)_\circ \xrightarrow{\mu} \mathrm{Aut}(p)_\circ$$

where μ is the multiplication (composition of maps) in $\mathrm{Aut}(p)_\circ$. Write

$$F * G\colon E \times S^p \times S^q \to E$$

for the adjoint map. Let $(\wedge(u, v)/\langle u^2, v^2\rangle, 0)$ denote the Sullivan model for $S^p \times S^q$ where $|u| = p$ and $|v| = q$.

Lemma 4.1. *A Sullivan model for the map $F * G$ is the DG algebra map*

$$\mathcal{A}_{F*G}\colon \wedge V \otimes \wedge W \to \wedge V \otimes \wedge W \otimes \wedge(u, v)/\langle u^2, v^2\rangle$$

given by

$$\mathcal{A}_{F*G}(\chi) = \chi + \theta_a(\chi)u + \theta_b(\chi)v + \theta_a \circ \theta_b(\chi)uv$$

for $\chi \in \wedge V \otimes \wedge W$.

Proof. The map $F * G$ is the composite

$$E \times S^p \times S^q \xrightarrow{F \times 1_{S^q}} E \times S^q \xrightarrow{G} E\,.$$

By the Künneth Theorem, a Sullivan model $\mathcal{A}_{F \times 1_{S^q}}$ for the product $F \times 1_{S^q}$ is the product of the Sullivan models $\mathcal{A}_F\colon \wedge V \otimes \wedge W \to \wedge V \otimes \wedge W \otimes (\wedge(u)/\langle u^2\rangle)$ and $1\colon \wedge(v)/\langle v^2\rangle \to \wedge(v)/\langle v^2\rangle$. Thus, given $\chi \in \wedge V \otimes \wedge W$ we see:

$$\begin{aligned}
\mathcal{A}_{F*G}(\chi) &= \mathcal{A}_{F \times 1_{S^q}}(\mathcal{A}_G(\chi)) \\
&= \mathcal{A}_{F \times 1_{S^q}}(\chi + \theta_b(\chi)v) \\
&= \mathcal{A}_F(\chi) + \mathcal{A}_F(\theta_b(\chi))v \\
&= \chi + \theta_a(\chi)u + \theta_b(\chi)v + \theta_a(\theta_b(\chi))uv.
\end{aligned}$$

□

We can now extend Theorem 3.3 to the fundamental group for the monoid $\mathrm{Aut}(p)_\circ$.

Theorem 4.2. *Let $p\colon E \to B$ be a fibration of simply connected CW complexes with E finite. Then the map*

$$\Phi'\colon \pi_1(\mathrm{Aut}(p)_\circ) \to H_1(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W))$$

is a homomorphism inducing an isomorphism

$$\Phi\colon \pi_1(\mathrm{Aut}(p)_\circ) \otimes \mathbb{Q} \xrightarrow{\cong} H_1(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W)).$$

Proof. To prove Φ' is a homomorphism, let $\alpha, \beta \in \pi_1(\mathrm{Aut}(p)_\circ))$ and recall the basic identity

$$\alpha \cdot \beta = (\alpha * \beta) \circ \Delta\colon S^1 \to \mathrm{Aut}(p)_\circ$$

where Δ is the diagonal map and the left-hand product is the usual multiplication in the fundamental group. Thus the adjoint to $\alpha \cdot \beta$ is the composition

$$S^1 \times E \xrightarrow{\Delta \times 1_E} S^1 \times S^1 \times E \xrightarrow{F*G} E$$

The result now follows directly from Lemma 4.1 and the fact that a Sullivan model for Δ is the map $\wedge(u, v) \to \wedge(w)$ given by $u, v \mapsto w$. The proof that Φ is a bijection is now the same as in the proof in Theorem 3.3. $\square$

We next prove the map $\Phi'\colon \pi_*(\mathrm{Aut}(p)_\circ) \to H_*(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W))$ preserves brackets which completes the proof of Theorem 1.

Theorem 4.3. *Let $p\colon E \to B$ be a fibration of simply connected CW complexes with E finite. Let $\alpha \in \pi_p(\mathrm{Aut}(p)_\circ)$ and $\beta \in \pi_q(\mathrm{Aut}(p)_\circ)$. Then*

$$\Phi'([\alpha, \beta]) = [\Phi'(\alpha), \Phi'(\beta)]\,,$$

where the left-hand bracket is the Samelson product in $\pi_*(\mathrm{Aut}(p)_\circ)$ and the right-hand bracket is that induced by the commutator in $\mathrm{Der}^*_{\wedge V}(\wedge V \otimes \wedge W)$.

Proof. By Proposition 2.2, $\mathrm{Aut}(p)_\circ$ is a grouplike space under composition. Let $\nu\colon \mathrm{Aut}(p)_\circ \to \mathrm{Aut}(p)_\circ$ be a homotopy inverse. Define $\overline{F}$ to be the composite

$$\overline{F}\colon E \times S^p \xrightarrow{1_E \times \alpha} E \times \mathrm{Aut}(p)_\circ \xrightarrow{1_E \times \nu} E \times \mathrm{Aut}(p)_\circ \xrightarrow{\omega} E$$

where ω is the evaluation map. Then $\overline{F}$ is adjoint to $-\alpha = \nu_\sharp(\alpha) \in \pi_p(\mathrm{Aut}(p)_\circ)$ and so has Sullivan model

$$\mathcal{A}_{\overline{F}}(\chi) = \chi - \theta_a(\chi)u\colon\ \wedge V \otimes \wedge W \to \wedge V \otimes \wedge W \otimes (\wedge(u)/\langle u^2\rangle)$$

for $\chi \in \wedge V \otimes \wedge W$ where $\langle \theta_a \rangle = \Phi'(\alpha)$.

Now given two classes $\alpha \in \pi_p(\mathrm{Aut}(p)_\circ)$ and $\beta \in \pi_q(\mathrm{Aut}(p)_\circ)$, the map Samelson product is defined by means of the map $\gamma\colon S^p \times S^q \to \mathrm{Aut}(p)_\circ$ defined by

$$\gamma(x, y) = \alpha(x) \circ \beta(y) \circ \overline{\alpha(x)} \circ \overline{\beta(y)}$$

where here $\overline{\alpha} = \nu_\sharp(\alpha)$. The map γ has adjoint Γ given by the composite

$$\begin{tikzcd}
E \times S^p \times S^q \arrow[r, "1_E \times \Delta \times \Delta"] \arrow[ddr, "\Gamma"'] & E \times S^p \times S^p \times S^q \times S^q \arrow[d, "1_E \times 1_{S^p} \times T \times 1_{S^q}"] \\
& E \times S^p \times S^q \times S^p \times S^q \arrow[d, "{[F,G]}"] \\
& E
\end{tikzcd}$$

where T is transposition and

$$[F, G] = F \circ (G \times 1_{S^n}) \circ (\overline{F} \times 1_{S^p \times S^q}) \circ (\overline{G} \times 1_{S^p \times S^q \times S^p})\,.$$

We see the map $[F, G]$ has Sullivan model

$$\mathcal{A}_{[F,G]}\colon\ \wedge V \otimes \wedge W \to \wedge V \otimes \wedge W \otimes (\wedge(u, v, \overline{u}, \overline{v})/\langle u^2, v^2, \overline{u}^2, \overline{v}^2\rangle)$$

given by

$$\begin{aligned}\mathcal{A}_{[F,G]}(\chi) \quad &= \chi + \theta_a(\chi)u + \theta_b(\chi)v - \theta_a(\chi)\overline{u} - \theta_b(\chi)\overline{v} \\ &\quad + \theta_a \circ \theta_b(\chi)uv + \theta_a \circ \theta_b(\chi)\overline{uv} - \theta_a \circ \theta_b(\chi)u\overline{v} - \theta_b \circ \theta_a(\chi)v\overline{u} \ . \\ &\quad + \text{ terms involving } u\overline{u} \text{ or } v\overline{v}\end{aligned}$$

Thus Γ has Sullivan model

$$\mathcal{A}_\Gamma(\chi) = \chi + [\theta_a, \theta_b](\chi)uv : \ \wedge V \otimes \wedge W \to \wedge V \otimes \wedge W \otimes \wedge(u,v)/\langle u^2, v^2\rangle$$

for $\chi \in \wedge V \otimes \wedge W$. The result now follows from the definition of the Samelson product: The restriction of Γ to $E \times (S^p \vee S^q)$ is null and so Γ induces $\Gamma' \colon E \times S^{p+q} \to E$ satisfying $\Gamma = \Gamma' \circ (1_E \times q)$ where $q \colon S^p \times S^q \to S^{p+q}$ is the projection onto the smash product. Finally, using the fact that q induces the map

$$w \mapsto uv : \ \wedge(w)/\langle w^2\rangle \to \wedge(u,v)/\langle u^2, v^2\rangle,$$

we see $\Phi'(\gamma)$ is represented by $[\theta_a, \theta_b]$. □

Remark 4.4. The techniques above can be applied to describe a related space of fibrewise self-homotopy equivalences. Let $F = p^{-1}(x)$ be the fibre over the base-point of $p \colon E \to B$ as above. The restriction res: $\mathrm{Aut}(p) \to \mathrm{Aut}(F)$ is a multiplicative continuous map. Denote the kernel by $\mathrm{Aut}^F(p)$. This monoid is of interest in the study of gauge groups. Denote by $\overline{\mathrm{Der}}_{\wedge V}(\wedge V \otimes \wedge W)$ the subcomplex of $\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge W)$ consisting of derivations θ such that $\theta(V) = 0$ and $\theta(W) \subset \wedge^+ V \otimes \wedge W$. Then we have a Lie algebra isomorphism

$$\pi_*(\mathrm{Aut}^F(p)_\circ) \otimes \mathbb{Q} \cong H_*(\overline{\mathrm{Der}}_{\wedge V}(\wedge V \otimes \wedge W)).$$

The details of the proof are extensive but entirely similar to the above.

5. Applications and Examples

We apply Theorem 1 to expand on some of the examples mentioned Section 2. We begin with the path-space fibration $p \colon PB \to B$ as discussed in Example 2.4. Let B be a simply connected CW complex. As in [10, Sec.16b], p has a relative minimal model of the form

$$I \colon (\wedge V, d) \to (\wedge V \otimes \wedge \overline{V}, D)$$

where, as usual, $(\wedge V, d)$ is the Sullivan minimal model for B. Here $\overline{V}$ is the desuspension of V, $(\overline{V})^n \cong V^{n-1}$. We recover part of the main result of [9].

Theorem 5.1. (Félix-Halperin-Thomas) *Let B be a simply connected CW complex and $p \colon PB \to B$ the path-space fibration. Then there is an isomorphism of graded Lie algebras*

$$\pi_*(\Omega B) \otimes \mathbb{Q} \cong H_*(\mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge \overline{V})).$$

Proof. The result is a direct consequence of Example 2.4 and Theorem 1. □

We next give a general bound for $\mathrm{Hnil}_{\mathbb{Q}}(\mathrm{Aut}(p)_\circ)$ in terms of the rational homotopy groups of the fibre.

Theorem 5.2. *Let $p \colon E \to B$ be a fibration of simply connected CW complexes with E finite. Let $F = p^{-1}(*)$ be the fibre. Then*

$$\mathrm{Hnil}_{\mathbb{Q}}(\mathrm{Aut}(p)_\circ) \le \mathrm{card}\{n \mid \pi_n(F) \otimes \mathbb{Q} \ne 0\}.$$

Proof. A nontrivial commutator $[\theta_1, \ldots, [\theta_{k-1}, \theta_k]]$ in $\mathrm{Der}^*_{\wedge V}(\wedge V \otimes \wedge W)$ gives rise to a sequence $w_1, \ldots, w_k$ of vectors in W with degrees $n_1 < \cdots < n_k$. To see this choose w_k so that $[\theta_1, \ldots, [\theta_{k-1}, \theta_k]](w_k) \neq 0$. Choose a nonvanishing summand $\theta_{i_1} \circ \cdots \circ \theta_{i_k}(w_k) \neq 0$. Choose w_{k-1} so that that w_{k-1} appears in $\theta_{i_k}(w_k)$ and $\theta_{i_1} \circ \cdots \circ \theta_{i_{k-1}}(w_{k-1}) \neq 0$. Proceed by induction. The result then follows from Theorem 1. □

For a principal G-bundle $p\colon E \to B$, we observe that $\mathrm{Hnil}_{\mathbb{Q}}(G)$ is often 1 (e.g., when G is a connected Lie group) while

$$\mathrm{Hnil}_{\mathbb{Q}}(\mathrm{Aut}(G)_\circ) = \mathrm{card}\{n \mid \pi_n(G) \otimes \mathbb{Q} \neq 0\}.$$

For recall G has minimal model $(\wedge(W), 0)$ with $W \cong \pi_*(G) \otimes \mathbb{Q}$ and so, in this case, $H_*(\mathrm{Der}(\wedge W); D) \cong \mathrm{Der}^*(\wedge W)$. Any sequence $w_{n_1}, \ldots, w_{n_k}$ of vectors in $W$of strictly increasing degree gives rise to a k-length commutator: Set $\theta_i(w_{n_i}) = w_{n_{i-1}}$ and $\theta_i(w_j) = 0$ where the w_j gives some homogeneous basis for W including the w_{n_i}. Thus we see

$$\mathrm{Hnil}_{\mathbb{Q}}(G_\circ) \leq \mathrm{Hnil}_{\mathbb{Q}}(\mathrm{Aut}(p)_\circ) \leq \mathrm{Hnil}_{\mathbb{Q}}(\mathrm{Aut}(G)_\circ).$$

Finally, we specialize to a case where we can make a complete calculation of the rational H-type of $\mathrm{Aut}(p)_\circ$.

Theorem 5.3. *Let $p\colon E \to B$ be a fibration with fibre $F = S^{2n+1}$ and E and B simply connected CW complexes with E finite. Suppose $i_\sharp\colon \pi_{2n+1}(S^{2n+1}) \to \pi_{2n+1}(E)$ is injective. Then $\mathrm{Aut}(p)_\circ$ is rationally homotopy abelian and, for each $q \geq 1$, we have isomorphisms*

$$\pi_q(\mathrm{Aut}(p)_\circ) \otimes \mathbb{Q} \cong H_{2n+1-q}(B; \mathbb{Q})\,.$$

Proof. Let $(\wedge V, d) \to (\wedge V \otimes \wedge(u), D)$ be a relative minimal model for p. A derivation $\theta \in \mathrm{Der}_{\wedge V}(\wedge V \otimes \wedge(u))$ is determined by the element $\theta(u) \in (\wedge V)^{2n+1-q}$. The derivation θ is a cocycle (resp. a coboundary) if $\theta(u)$ is a cocycle (resp. a coboundary). Further, the commutator bracket of any two such derivations is directly seen to be trivial. The result thus follows from Theorem 1. □

6. On the group of components of $\mathrm{Aut}(p)$

As mentioned in the introduction, the group $\mathcal{E}(p) = \pi_0(\mathrm{Aut}(p))$ of path components of $\mathrm{Aut}(p)$ does not generally localize well. First $\mathcal{E}(p)$ is often non-nilpotent. Even when $\mathcal{E}(p)$ is nilpotent, it may not satisfy $\mathcal{E}(p)_{\mathbb{Q}} = \mathcal{E}(p_{\mathbb{Q}})$ — take $p\colon S^n \to *$, for an easy example. However, by work of Dror-Zabrodsky [8] and Maruyama (e.g., [23]), various natural subgroups of $\mathcal{E}(p)$ are nilpotent and do localize well. We consider one such example which we denote $\mathcal{E}_\sharp(p)$. We note that different versions of groups of fibrewise equivalences have also been considered [13, 11]. Our definition of $\mathcal{E}_\sharp(p)$ is chosen to allow for an identification of $(\mathcal{E}_\sharp(p))_{\mathbb{Q}}$ in the spirit of Theorem 1.

Suppose, as usual, that $p\colon E \to B$ is a fibration of simply connected CW complexes with E finite. Then any fibrewise self-map of E is fibrewise homotopic to a based self-map. Further, any two based fibrewise self-maps of E that are freely fibrewise homotopic are also based fibrewise homotopic. Thus in what follows we work with based fibre-self-equivalences of $p\colon E \to B$.

Consider the long exact sequence of the fibration

$$\begin{tikzcd} \cdots \arrow[r] & \pi_{i+1}(B) \arrow[r, "\partial"] & \pi_i(F) \arrow[r, "j_\#"] & \pi_i(E) \arrow[r, "p_\#"] & \pi_i(B) \arrow[r] & \cdots \end{tikzcd}$$

Each $f \in \mathrm{Aut}(p)$ induces an automorphism $f_\#$ of this sequence, which is the identity on $\pi_*(B)$. Thus $f_\#$ induces an automorphism of cokernels

$$\overline{f_\#} : \frac{\pi_*(F)}{\mathrm{im}\partial} \to \frac{\pi_*(F)}{\mathrm{im}\partial}.$$

Define $\mathcal{E}_\#(p)$ to be the group of based homotopy classes of based equivalences in $\mathrm{Aut}(p)$ that induce the identity on $\mathrm{im}\partial \subseteq \pi_*(F)$ through degree equal to the dimension of E, and also induce the identity on the cokernels above through the dimension of E. Notice that this reduces to the subgroup of classes that induce the identity on $\pi_*(F)$ through degree equal to the dimension of E if the fibration is "Whitehead trivial," i.e., if the connecting homomorphism ∂ is trivial (through the dimension of E) as considered in [11]. When p is trivial, it reduces to the subgroup $\mathcal{E}_\sharp(E)$ of classes that induce the identity on $\pi_*(E)$ through degree equal to the dimension of E as in [23].

Theorem 6.1. *Let $p\colon E \to B$ be a fibration of simply connected CW complexes with E finite. Then $\mathcal{E}_\sharp(p)$ is a nilpotent group. Given any set of primes $\mathbb{P}$ we have*

$$\mathcal{E}_\#(p)_{\mathbb{P}} \cong \mathcal{E}_\#(p_{\mathbb{P}}).$$

Proof. Observe $\mathcal{E}_\sharp(p)$ acts nilpotently on the normal chain

$$0 \triangleleft \mathrm{im}\partial \triangleleft \pi_*(F)$$

through the dimension of E. That $\mathcal{E}_\sharp(p)$ is a nilpotent group follows from [8]. The result on localization is now obtained by adjusting Maruyama's argument [23] to the fibrewise setting by replacing the use of a Postnikov decomposition of E with Moore-Postnikov decomposition of $p\colon E \to B$. We omit the details. □

By virtue of Theorem 6.1, we may identify $\mathcal{E}_\#(p)_{\mathbb{Q}}$ in terms of certain automorphisms of the Sullivan model of E. As usual, let $(\wedge V, d) \to (\wedge V \otimes \wedge W, D)$ denote the minimal model of p. Let $D_0 : W \to V$ be the linear part of D, and decompose W as $W = W_0 \oplus W_1$ with $W_0 = \mathrm{ker} D_0$ and W_1 a complement. Topologically, W_1 may be identified with $\mathrm{im}\partial = \mathrm{ker} j_\sharp$ and W_0 with $\mathrm{coker}\partial = \mathrm{im} j_\sharp$ after rationalization. Write $\mathrm{Aut}_\sharp(\wedge V \otimes \wedge W)$ for the group of automorphisms φ of $\wedge V \otimes W$ which are the identity on $\wedge V$ and have linear part $\varphi_0 : W \to V \oplus W$ satisfying $(\varphi_0 - 1)(W_0) \subseteq V$ and $(\varphi_0 - 1)(W_1) \subseteq V \oplus W_0$. As in [2], we may identify $\mathcal{E}_\#(p_{\mathbb{Q}})$ with the group of homotopy classes of $\mathrm{Aut}_\sharp(\wedge V \otimes \wedge W)$.

Recall from the introduction that $\mathrm{Der}^0_\#(\wedge V \otimes \wedge W)$ denotes the vector space of derivations θ of degree 0 of $\wedge V \otimes \wedge W$ that satisfy $\theta(V) = 0$, $\mathcal{D}(\theta) = 0$, $\theta(W_0) \subset V \oplus \wedge^{\geq 2}(V \oplus W)$, and $\theta(W_1) \subset V \oplus W_0 \oplus \wedge^{\geq 2}(V \oplus W)$. The derivation differential defines a linear map

$$\mathcal{D} : \mathrm{Der}^1_{\wedge V}(\wedge V \otimes \wedge W) \to \mathrm{Der}^0_\#(\wedge V \otimes \wedge W)\,, \qquad \theta \mapsto \mathcal{D}(\theta) = D\theta + \theta D.$$

We are writing

$$H_0(\mathrm{Der}_\sharp(\wedge V \otimes \wedge W)) = \mathrm{coker}\,\mathcal{D}\,.$$

As in [29, Sec.11] and [26, Pro.12], the correspondence $\theta \mapsto e^\theta$ gives a bijection (with inverse $\varphi \mapsto \log(\varphi)$) from $\mathrm{Der}^0_\#(\wedge V \otimes \wedge W)$ to $\mathrm{Aut}_\sharp(\wedge V \otimes \wedge W)$. Under this bijection, compositions of automorphisms $e^\theta \circ e^\phi$ correspond to "Baker-Campbell-Hausdorff" products of derivations, i.e.,

$$\log(e^\theta \circ e^\phi) = \theta + \phi + \frac{1}{2}[\theta, \phi] + \frac{1}{12}[\theta, [\theta, \phi]] + \cdots .$$

Theorem 6.2. *Let $p\colon E \to B$ be a fibration of simply connected CW complexes with E finite. The assigment $\theta \mapsto e^{\theta}$ induces an isomorphism of groups*

$$\Psi\colon H_0(\mathrm{Der}_\sharp(\wedge V \otimes \wedge W)) \to \mathcal{E}_\#(p_{\mathbb{Q}}),$$

where Baker-Campbell-Hausdorff composition of derivations is understood in the left-hand term.

Proof. To see that Ψ is well-defined, we observe that the assignment $\theta \mapsto e^{\theta}$ restricts to a correspondence between boundaries in $\mathrm{Der}^0_\sharp(\wedge V \otimes \wedge W)$ and automorphisms in $\mathrm{Aut}_\sharp(\wedge V \otimes \wedge W)$ homotopic to the identity there. Suppose $\theta = \mathcal{D}(\theta_1)$ with $\theta_1 \in \mathrm{Der}^1_{\wedge V}(\wedge V \otimes \wedge W)$. Then we define a derivation θ_2 of $\wedge V \otimes \wedge W \otimes \wedge(t, dt)$ by $\theta_2(V) = \theta_2(t) = \theta_2(dt) = 0$ and $\theta_2(w) = t\theta_1(w)$ for $w \in W$. The map

$$\mathcal{H} = e^{\mathcal{D}(\theta_2)} : \wedge V \otimes \wedge W \to \wedge V \otimes \wedge W \otimes \wedge(t, dt)$$

satisfies $p_1 \circ \mathcal{H} = e^{\theta}$ and $p_0 \circ \mathcal{H}$ is the identity. Conversely, a given homotopy $\mathcal{H}\colon \wedge V \otimes \wedge W \to \wedge V \otimes \wedge W \otimes \wedge(t, dt)$ between the identity and φ in $\mathrm{Aut}_\sharp(\wedge V \otimes \wedge W)$ may be chosen to be fibrewise in the sense of Proposition 3.2. Taking the dt component of $\log(\mathcal{H})$ we obtain a derivation $\theta \in \mathrm{Der}^1_{\wedge V}(\wedge V \otimes \wedge W)$ with $\mathcal{D}(\theta) = \log(\varphi)$. The proof that the induced map Ψ is a bijection now follows the same line as the proof of Theorem 2. $\square$

We remark that this result may be used to analyze the nilpotency of $\mathcal{E}_\#(p_{\mathbb{Q}})$, and ultimately, since we have $\mathcal{E}_\#(p_{\mathbb{Q}}) \cong \mathcal{E}_\#(p)_{\mathbb{Q}}$, that of $\mathcal{E}_\#(p)$, in terms of bracket lengths in $H_0(\mathrm{Der}_\#(\wedge V \otimes \wedge W))$.

Finally, observe that the results of this section may also be applied to give cardinality results for the order of $\mathcal{E}(p)$. As a simple example, let $\pi\colon B \times S^n \to B$ be a trivial fibration. If $H^n(B; \mathbb{Q}) = 0$ then $\mathcal{E}(\pi)$ is a finite group. On the other hand denote by V^n the dual of the image of the rational Hurewicz map in $H^n(B; \mathbb{Q})$. Then clearly $\mathcal{E}_\#(\pi_{\mathbb{Q}}) \cong H_0(\mathrm{Der}_\sharp(\wedge V \otimes \wedge W)) \cong V^n$. Thus if V^n is nontrivial then $\mathcal{E}(\pi)$ is infinite in this case. We conclude with one example involving a nontrivial fibration.

Example 6.3. Let $p\colon S^7 \times S^3 \to S^4$ be the composition $p = \eta \circ p_1$, where η denotes the Hopf map and p_1 projection onto the first factor. Then the fibre is $S^3 \times S^3$. A relative minimal model for p is given by

$$\wedge(v_4, v_7) \to \wedge(v_4, v_7) \otimes \wedge(w_3, w_3') \to \wedge(w_3, w_3'),$$

with (non-minimal) differential given by $D(v_7) = v_4^2$, $D(w_3') = v_4$, and $D = 0$ on other generators. With reference to our notation above, we have $V = \langle v_4, v_7 \rangle$, $W_0 = \langle w_3 \rangle$, and $W_1 = \langle w_3' \rangle$. Define a differential $\theta \in \mathrm{Der}^0_\#(\wedge V \otimes \wedge W)$ by setting $\theta(w_3') = w_3$, and $\theta = 0$ on other generators. A direct check shows that θ represents a non-zero class in $H_0(\mathrm{Der}_\sharp(\wedge V \otimes \wedge W))$. From Theorem 6.3 and the discussion of this section, it follows that $\mathcal{E}_\#(p)$ has infinite order.

Under the isomorphism of Theorem 6.3, the derivation θ evidently corresponds to the automorphism φ of $\wedge V \otimes \wedge W$ given by $\varphi(w_3') = w_3' + w_3$ and $\varphi = 1$ on other generators. Notice that this automorphism does not correspond to an element of $\mathcal{E}_\#(F)$—in the "non-fibrewise" notation of [8]. Also, since in this case the total space has the rational homotopy type of $S^3 \times S^7$, we see that φ does not correspond to an element of $\mathcal{E}_\#(E)$, which is trivial rationally. The example thus demonstrates that, in general, it is necessary to consider the relative minimal model of a fibration and not just the minimal models of the spaces invloved to analyze $\mathcal{E}_\#(p)$ rationally.

References

[1] M. Arkowitz and C. R. Curjel, *Homotopy commutators of finite order. II.* Quart. J. Math. Oxford Ser. (2) **15** 1964 316–326

[2] M. Arkowitz and G. Lupton, *On the nilpotency of subgroups of self-homotopy equivalences*, Algebraic topology: new trends in localization and periodicity, Progr. Math., **136**, 1–22.

[3] I. Berstein and T. Ganea, *Homotopical nilpotency*, Illinois J. Math. **5** (1961), 99–130.

[4] J. Block and A. Lazarev *André-Quillen cohomology and rational homotopy of function spaces,* Adv. Math. **193** (2005), no. 1, 18–39.

[5] P. Booth, P. Heath, C. Morgan, R. Piccinini *H-spaces of self-equivalences of fibrations and bundles,* Proc. London Math. Soc. (3) **49** (1984), no. 1, 111–127.

[6] U. Buijs and A. Murillo, *The rational homotopy Lie algebra of function spaces*, Comment. Math. Helv. 83 (2008), no. 4, 723–739.

[7] M. C. Crabb and I. M. James, *Fibrewise homotopy theory*, Springer-Verlag, 1998.

[8] E. Dror and A. Zabrodsky *Unipotency and nilpotency in homotopy equivalences,* Topology **18** (1979), no. 3, 187–197.

[9] Y. Félix, S. Halperin and J.-C. Thomas, *Sur certaines algébres de Lie de dérivations*, Annales Institut Fourier 32 (1982), 143-150.

[10] ———, *Rational homotopy theory*, Springer-Verlag, 2001.

[11] Y. Félix and A. Murillo, *A note on the nilpotency of subgroups of self-homotopy equivalences,* Bull. London Math. Soc. **29** (1997),486–488.

[12] Y. Félix and J. Oprea, *Rational homotopy of gauge groups,* to appear in Proc. Amer. Math. Soc.

[13] Y. Félix and J.-C. Thomas, *Nilpotent subgroups of the group of fibre homotopy equivalences,* Publ. Mat. **39** (1995), 95–106

[14] S. Halperin, *Finiteness in the minimal models of Sullivan,* Trans. Amer. Math. Soc. **230** (1977), 173-199.

[15] M. Fuchs, *Verallgemeinerte Homotopie-Homomorphismen und klassifizierende Räume*, Math. Ann. **161** (1965), 197–230.

[16] D. Gottlieb, *On fibre spaces and the evaluation map*, Ann. Math. **87** (1968), 42–55.

[17] P. Hilton, G. Mislin, and J. Roitberg, *Localization of nilpotent groups and spaces*, North-Holland Publishing Co., Amsterdam, 1975, North-Holland Mathematics Studies, No. 15, Notas de Matemática, No. 55 [Notes on Mathematics, No. 55].

[18] P. J. Kahn, *Some function spaces of CW type*, Proc. Amer. Math. Soc. **90**, 599-607.

[19] J. R. Klein, C. L. Schochet and S. B. Smith, *Continuous trace C^*-algebras, gauge groups and rationalization,* preprint, arXiv:0811.0771v1.pdf

[20] G. Lupton, N. C. Phillips, C. L. Schochet and S. B. Smith, *Banach algebras and rational homotopy theory*, Trans. Amer. Math. Soc. **361** (2009), 267-295.

[21] G. Lupton and S. B. Smith, *Rationalized evaluation subgroups of a map I: Sullivan models, derivations and G-sequences*, J. Pure and Applied Algebra 209 (2007), 159-171.

[22] ———, *Rank of the fundamental group of any component of a function space*, Proc. Amer. Math. Soc. 135 (2007), 2649-2659.

[23] K. Maruyama, *Localization of a certain subgroup of self-homotopy equivalences*, Pacific J. Math. 136 (1989), 293–301

[24] W. Meier, *Rational universal fibrations and flag manifolds,* Math. Ann. **258** (1982), 329-340.

[25] J. Milnor, *On spaces having the homotopy type of CW-complex*, Trans. Amer. Math. Soc. **90** (1959), 272-280.

[26] P. Salvatore, *Rational homotopical nilpotency of self-equivalences,* Topology and its Appl., **77** (1997), 37- 50.

[27] H. Scheerer, *On rationalized H- and co-H-spaces. With an appendix on decomposable H- and co-H-spaces*, Manuscripta Math. **51** (1985), 63–87. MR 88k:55007

[28] H. Shiga and M. Tezuka, *Rational fibrations, Homogeneous spaces with positive Euler characteristic and Jacobians*, Ann. Inst. Fourier 37 (1987), 81-106.

[29] D. Sullivan, *Infinitesimal comutations in topology*, Publ. I.H.E.S. **47** (1977), 269-331.

[30] G. Whitehead, *Elements of homotopy theory*, Graduate Texts in Mathematics, vol. 61, Springer-Verlag, New York, 1978.

Institut Mathématique, Université Catholique de Louvain, B-1348 Louvain-la-Neuve Belgique
E-mail address: `felix@math.ucl.ac.be`

Department of Mathematics, Cleveland State University, Cleveland OH 44115 U.S.A.
E-mail address: `G.Lupton@csuohio.edu`

Department of Mathematics, Saint Joseph's University, Philadelphia, PA 19131 U.S.A.
E-mail address: `smith@sju.edu`